

Glossary

Algae. Small aquatic plants containing chlorophyll and without roots that occur as a single cell or multi-celled colonies. Algae form the base of the food chain in an aquatic environment.

Algal bloom. A heavy growth of algae in and on a body of water as a result of high nutrient concentrations.

Aquatic Invasive Species (AIS). Refers to species of plant or animal that are not native to a particular region into which they have moved or invaded. Zebra mussels and Eurasian water-milfoil are examples of AIS. Wisconsin has laws preventing the spread on boats and trailers.

Bathymetric map. A map showing depth contours in a water body. Bottom contours are usually presented as lines of equal depth, in meters or feet. Often called a hydrographic map.

Chlorophyll. Green pigment present in all plant life and necessary for photosynthesis. The amount of chlorophyll present in lake water depends on the amount of algae and is used as a common indicator of water quality.

Cultural eutrophication. Accelerated eutrophication of a lake that occurs as a result of human activities in the watershed. These activities increase nutrient loads in runoff water that drains into lakes.

Decomposition. The act of breaking down organic matter from a complex form to a simpler form, mainly through the action of fungi and bacteria.

Deionized water. Water that has been passed through a column or membrane to remove ions present.

Distilled water. Water that is boiled in a still and the condensate collected and distributed. Distillation removes both ionic and nonionic organic contaminants.

Dissolved oxygen. A measure of the amount of oxygen gas dissolved in water and available for use by microorganisms and fish. Dissolved oxygen is produced by aquatic plants and algae as part of photosynthesis.

Drainage lake. Lakes fed primarily by streams and with outlets into streams or rivers. They are more subject to surface runoff problems but generally have shorter residence times than seepage lakes. Watershed protection is usually needed to manage lake water quality.

Epilimnion. The uppermost circulating layer of warm water that occurs in stratified lakes in summer because of the differences in water density.

Euphotic zone. That part of a water body where light penetration is sufficient to maintain photosynthesis.

Eutrophic. Lakes characterized by high nutrient inputs, high productivity, often experiencing algal blooms and abundant weed growth. This term can also refer to a nutrient-rich lake, as large amounts of algae and weeds characterize a eutrophic lake.

Eutrophication. The process by which lakes and streams are enriched by nutrients causing an increase in plant and algae growth.

Georegion. Wisconsin's lake "georegions" originated from a grouping of lakes made in the early 1980s by Wisconsin DNR senior limnologists. These groupings are based on the best professional judgment of the scientists most familiar with Wisconsin's lake resources. The georegions roughly reflect "hydro-chemical lake regions" which are based on the state's bedrock geology, glacial geology and soil type, and the more recently described "ecoregions" which are based on geological characteristics as well as the dominant vegetation.

Hypolimnion. The cold, deepest layer of a lake that is removed from surface influences.

(Glossary continued on next page)

Lake association. A voluntary organization with a membership generally comprised of those who own land on or near a lake. The goals of lake associations usually include maintaining, protecting, and improving the quality of a lake, its fisheries, and its watershed.

Lake classification. A way of placing lakes into categories with management strategies best suited to the types of lakes found in each category. For example, lakes can be classified to apply varying shoreland development standards. They can be grouped based on hydrology, average depth, surface area, shoreline configuration, as well as, sensitivity to pollutants and recreational use.

Lake district. A special purpose unit of government with the cause of maintaining, protecting, and improving the quality of a lake and its watershed for the mutual good of the members and the lake environment.


Limnology. The study of inland lakes and waters. The study of the interactions of the biological, chemical, and physical parameters of lakes and rivers.

Macrophyte. Large, rooted or floating aquatic plants that may bear flowers and seeds.

Meniscus. The curved upper surface of a still liquid in a tube caused by surface tension.

Mesotrophic. Lakes characterized by their moderately fertile nutrient levels. Falls in between the oligotrophic and eutrophic levels of nutrient enrichment.

Metalimnion. Sometimes referred to as the thermocline. The narrow transition zone between the epilimnion and the hypolimnion that occurs in stratified lakes.



Nitrogen. One of the major nutrients required for the growth of aquatic plants and algae.

Oligotrophic. Lakes characterized by low nutrient inputs and low productivity. They are generally deep with high water clarity.

Parts per million (ppm). An expression of concentration indicating weight of a substance in a volume of one liter. Milligrams per liter (mg/l) is an equivalent unit.

pH. The measure of the acidity or alkalinity of a solution. Neutral solutions are defined as having a pH of 7.0. Solutions which are known as acidic have a pH lower than 7. Solutions which are known as basic have a pH greater than 7.

Phosphorus. The major nutrient influencing plant and algal growth in more than 80% of Wisconsin lakes. Soluble reactive phosphorus refers to the amount of phosphorus in solution that is available to plants and algae. Total phosphorus refers to the amount of phosphorus in solution (reactive) and in particulate forms (non-reactive.)

Photic zone. The surface and underwater lighted zone in a lake that usually has a depth around 1.7 times the Secchi reading.

Photosynthesis. Process by which green plants convert carbon dioxide (CO₂) dissolved in water to sugar and oxygen using sunlight for energy. Photosynthesis is essential in producing a lake's food base and is an important source of oxygen for many lakes.

Phytoplankton. Very small free-floating aquatic plants, such as one-celled algae. Their abundance, as measured by the amount of chlorophyll a in a water sample, is commonly used to classify the trophic status of a lake.

Qualified Lake Association. To be eligible for state lake planning, protection and recreational boating facilities grants, a lake association must meet certain standards set out in section 281.68 of the Wisconsin statutes.

(Glossary continued on next page)

Respiration. The reverse reaction of photosynthesis. The complex process that occurs in the cells of plants and animals in which nutrient organic molecules, such as glucose, combine with oxygen to produce carbon dioxide, water, and energy. Respiration consumes oxygen and releases carbon dioxide. This process also takes place during decomposition as bacterial respiration increases.

Runoff. Water from rain, snow melt, or irrigation that flows over the ground surface and into streams or lakes.

Secchi disk. A 20-cm (8-inch) diameter disk painted white and black in alternating quadrants. It is used to measure light transparency in lakes.

Seepage lakes. Lakes without a significant inlet or outlet, fed by rainfall and groundwater.

Spring lakes. Lakes that have no inlet, but have an outlet. The primary source of water for spring lakes is groundwater flowing into the bottom of the lake from inside and outside the immediate surface drainage area. Spring lakes are found at the headwaters of many streams and are a fairly common type of lake in northern Wisconsin.

State Laboratory of Hygiene. The state of Wisconsin's public health and environmental laboratory.

Station # (or Storet #). A number assigned to sampling locations on a waterbody. The Station # makes it easy to track secchi and chemistry data. Each sampling site on a lake will have a separate Station #.

Stratification. The layering of water due to differences in temperature and density.

SWIMS. Surface Water Integrated Monitoring System. The database where all CLMN data and other water quality data is stored.

Tannins. Natural pigments found in organic matter such as leaves and bark.

Thermocline. Sometimes referred to as the metalimnion. The narrow transition zone between the epilimnion and the hypolimnion that occurs in stratified lakes.

Trophic state. The extent to which the process of eutrophication has occurred is reflected in a lake's trophic classification or state. The three major trophic states are oligotrophic, mesotrophic, and eutrophic.

Turion. A specialized bud which consists of condensed leaves and stems. This structure is most often an "over-wintering" structure, but in the case of curly-leaf pondweed is an "over-summering" structure. When the appropriate water conditions are reached, the turion will sprout a new plant.

µg/L. micrograms per liter is an expression of concentration indicating weight of a substance in a volume of one liter. Parts per billion (ppb) is an equivalent unit.

Waterbody # or WBIC (Waterbody Identification Code).

A unique identification number the Wisconsin DNR uses to identify each waterbody in the state. Every one of the 15,000 lakes in Wisconsin has a unique WBIC.

Watershed. The area of land draining into a specific stream, river, lake, or other body of water.

Zebra mussel. A tiny bottom dwelling mollusk native to Europe.

Zooplankton. Plankton that is made up of microscopic animals, for example, protozoa, that eat algae. These suspended plankton are an important component of the lake food chain and ecosystem. For many fish and crustaceans, they are the primary source of food.